

- machulkai* (*Sternocera castanea rothschildi* ab.), Obenberger, 1935: 99.
«Typus» (7221), Kenia: Laikipia, Uaso Nyiro, Afr. O. (Machulka).
- melichari* (*Kheilila*), Obenberger, 1923, *Zeitschr. Wiss. Ins. Biol.*, II: 114.
Syntypus (7241), Perù: Puno (Soukup).
- minor* (*Paragrilus paulensis* var.), Obenberger, 1924a: 143. «Typus» (7254), Brasile: Sao Paulo (J. Mráz).
- mrazi* (*Taphrocerus*), Obenberger, 1924b: 63. «Typus» (7219), Brasile: Sao Paulo (J. Mráz).
- olsoufieviella* (*Polybothris*), Obenberger, 1942: 32. «Typus» (7245), Madagascar (coll. Nonfried).
- paulensis* (*Paragrilus*), Obenberger, 1924a: 142. «Typus» (7239), Brasile: Sao Paulo (J. Mráz).
- pragensis* (*Agrilus cyaneus* ab.), Obenberger, 1917: 214. «Typus» (7223), [Cecoslovacchia]: V. Osek, Boh[emia] (J. Steigervald).
- preyssleri* (*Agrilus*), Obenberger, 1933a: 19. «Typus» (7231), [Brasile]: Sao Paulo (J. Mráz).
- pseudosepulchralis* (*Anthaxia nigrojubata* var.), Obenberger, 1938: 227. «Typus» (7227), Grecia: [M.] Olympos, m 2100, 4.VI.1937, (coll. Barton).
- punctipennis* (*Aphanisticus*), Kerremans, 1906: 102. Holotypus (7229), [Etiopia]: Eritrea, Adi-Ugri, VII.1901 (A. Andreini).
- purpurescens* (*Chrysobothris*), Kerremans, 1906: 98. Holotypus (7243), [Etiopia]: Eritrea, Asmara, Arbaroba (A. Andreini).
- rutiliplaga* (*Polybothris sumptuosa* var.), Obenberger, 1928: 167. «Typus» (7249), Madagascar.
- sekerai* (*Agrilus*), Obenberger, 1933a: 13. «Typus» (7232), Brasile: Sao Paulo (J. Mráz).
- spesivcevi* (*Polybothris*), Obenberger, 1928: 174. «Typus» (7247), Madagascar.
- staudingerina* (*Lampetis*), Obenberger, 1917: 278. «Typus» (7251), Madagascar.
- subnodosa* (*Trachys*), Kerremans, 1906: 103. Holotypus (7217), [Etiopia]: Eritrea, Adi-Ugri, Mai Daro (A. Andreini).
- ugriana* (*Sphenoptera*), Kerremans, 1906: 94. Holotypus (7216), [Etiopia]: Eritrea, Adi-Ugri (A. Andreini).
- zoufali* (*Polybothris*), Obenberger, 1928: 171. «Typus» (7250), Madagascar.
- xenia* (*Damarsila*), Obenberger, 1924a: 57. «Typus» (7236) «D. O. Afr.»: Manow.

ELATEROIDEA

ELATERIDAE

- aethiopicus* (*Propsephus*), Girard, 1976: 424. Holotypus (7181), Africa Orientale [ex] italiana: tra Bongo e Uota, 17.III.1939 (L. Cipriani).
- brachypilus* (*Pseudolophoeus*), Girard, 1984: *Ann. Soc. entomol. France*, (n.s.) 3 (in stampa). Holotypus (3080), [Etiopia]: Borana, Javello, XI.1938-III.1939 (E. Faà di Bruno).
- ineditus* (*Propsephus*), Girard, 1976: 427. Holotypus (3075), [Etiopia]: Borana, Javello, XI.1938-III.1939 (E. Faà di Bruno).
- pilosus* (*Anisomerus*), Girard, 1984: *Revue fr. Entomol.*, (n.s.) 3 (in stampa). Holotypus (3063), Somalia: Jesomma, 15.VIII.1968 (S.B.S.), (*). Paratypi (3064), stessi dati, 3 ex., (*). Paratypus (3065), Somalia: Bud-Bud, 15.VIII.1968 (S.B.S.), (*).
- proximus* (*Propsephus*), Girard, 1984: *Ann. Soc. entomol. France*, (n.s.) 3 (in stampa). Holotypus (3077), Somalia: Gelib, Alessandra, IV.1937 (F. Bigi). Paratypus (3078), stessi dati.
- zairensis* (*Eudicronychus*), Girard, 1948: *Revue franc. Entomol.*, (n.s.) 3 (in stampa). Paratypi (6787), Zaïre: Buie, 27.III-5.IV.1975 (S.B.Z.), 33 ex., (*).

CANTHAROIDEA

LAMPYRIDAE

- flavocostata* (*Luciola bimaculicollis* var.), Pic, 1928a: 21. Holotypus (7141), Somalia [ex] italiana: Margherita, VI.1913.
- latevittata* (*Luciola gaudens* var.), Pic, 1928a: 21. Holotypus (7142), Somalia [ex] italiana: Margherita, VI.1913.

BOSTRYCHOIDEA

PTINIDAE

- chiaromonte* (*Ptinus*), Pic, 1928a: 39. Holotypus (7166), Somalia [ex] italiana meridionale: Vill. Duca Abruzzi, III-IV.1926 (Miss. ent. Paoli).

MELYRIDAE
(inclusi MALACHIDAE)

- minutemaculatus* (*Pseudocoletes*), Wittmer, 1978: 100. Holotypus (6081), Somalia: Sar Uanle, 6.XI.1971 (S.B.S.-P.L.), (*). Paratypi (6078, 6079), stessa località, 3-10.XI.1971 e 28.V.1973 (S.B.S.-P.L.), 17 ex, (*).
- paolii* (*Mixis*), Pic, 1928b: 37. Holotypus (7138), Somalia [ex] italiana meridionale: Vill. Duca Abruzzi, 1926 (Miss. ent. Paoli). Paratypi (7139), Somalia [ex] italiana meridionale: Vill. Duca Abruzzi, IX.1926 (Miss. ent. Paoli), 3 ex.
- testaceipes* (*Pseudocoletes*), Pic, 1928b: 38. Holotypus (7140), Somalia [ex] italiana meridionale: Vill. Duca Abruzzi, X.1926 (Miss. ent. Paoli).

CUCUJOIDEA

COCCINELLIDAE

- erithraeae* (*Alesia trilineata* ssp.), Korschefsky, 1940: 72. Syntypi (7170), [Etiopia]: Eritrea, Adi-Ugri, VII.1901 (A. Andreini). 13 ex. Syntypus (7171), [Etiopia]: Eritrea, Conca di Bahat, 20.IX.1901 (A. Andreini).
- martellii* (*Nephus*), Fürsch, 1979: 21. Holotypus ♀ (2065), [Etiopia]. Eritrea, Adi-Caié, VIII.1902 (A. Andreini).
- somaliensis* (*Nephus*), Fürsch, 1978: 266. Holotypus (2063), Somalia: Sar Uanle, 13.VI.1973 (S.B.S.-P.L.), (*). Allotypus (3501), stessa località, 14.VI.1973 (S.B.S.-P.L.), (*). Paratypi (2064, 3502, 3503), stessa località, 17.V-8.VI.1973 (S.B.S.-P.L.), 3 ex, (*).
- zimmermanni* (*Solanophila*), Weise, 1922: 105. Syntypus (7172), [Tanzania]: Amani (Zimmermann).

TENEBRIONIDAE

- andreinii* (*Catomus*), Gridelli, 1940: 253. Holotypus (7191), [Etiopia]: Eritrea, Adi-Caié, IX.1902 (A. Andreini).
- andreinii* (*Gonocnemis*), Gridelli, 1940: 251. Holotypus ♀ (7190), [Etiopia]: Eritrea, Adi-Ugri, VIII.1901 (A. Andreini).
- andreinii* (*Hypophloeus*), Gridelli, 1940: 246. Paratypus (7188), [Etiopia]: Eritrea, Adi-Caié, 20.IV.1902 (A. Andreini).

- andreinii* (*Oxycara aethiopicum* ssp.), Gridelli, 1940: 222. Holotypus (7186), [Etiopia]: Eritrea, Saganeiti, IV.1901 (A. Andreini). Paratypi (7187), stessi dati, 5 ex.
- andreinii* (*Pachychilina*), Doderò, 1916: 350. Syntypi (7185), [Libia]: Tripolitania, Misurata, X.1913 (A. Andreini), 4 ex.
- andreinii* (*Tenebrio*), Gridelli, 1940: 247. Syntypi (7192), [Etiopia]: Eritrea, Adi-Ugri, IX.1901 (A. Andreini), 4 ex.
- carfi* (*Stenosis brenthoides* ssp.), Ardoin, 1971: 46. Holotypus ♂ (129), Tunisia: Isola Gallo, arc. Galita, 2.VI.1966 (B. Lanza e S. Carfi). Allotypus ♀ (130), stessi dati. Paratypi (131), stessi dati, 6 ex.
- dubia* (*Oxycara aethiopicum* ssp.), Gridelli, 1940: 223. Holotypus (7203), Etiopia: Dongollo, 29-30.XII.1900 (A. Andreini).
- erythraeus* (*Hypophloeus*), Gridelli, 1940: 245. Paratypi (7189), [Etiopia]: Eritrea, Adi-Caié, 20.IV.1902 (A. Andreini), 2 ex.
- erythraeus* (*Opatrinus corvinus* ssp.), Gridelli, 1940: 239. Syntypus (7210) [Etiopia]: Eritrea, Adi-Ugri, 16-30.VI.1901 (A. Andreini).
- griseovittatum* (*Gonocephalum*), Gridelli, 1940: 241. Syntypi (7193), [Etiopia]: Eritrea, Adi-Caié, V.1902 (A. Andreini), 4 ex. Syntypi (7194), [Etiopia]: Eritrea, Conca di Bahat, 16-21.IX.1901 (A. Andreini), 2 ex. Syntypus (7196), [Etiopia]: Eritrea, Adi-Caié, VII.1902 (A. Andreini). Syntypus (7197), [Etiopia]: Eritrea, Ghinda, 11.XI.1901 (A. Andreini). Syntypus (7195), [Etiopia]: Eritrea, Adi-Ugri, 16-30.VI.1901 e V.1901 (A. Andreini), 2 ex.
- inhumeralis* (*Praogena*), Pic, 1928b: 43. Holotypus (7207), Somalia mer.: Villaggio Duca Abruzzi, 1926 (Miss. ent. Paoli).
- lanzai* (*Opatrum*), Ardoin, 1971: 51. Holotypus ♂ (147), Tunisia: Isola Galitone, arc. Galita, 30.V.1966 (B. Lanza e S. Carfi). Allotypus ♀ (148), stessi dati. Paratypi (149), stessi dati, 5 ex.
- laticollis* (*Praogena*), Pic, 1928a: 25. Holotypus (7209), Somalia: Lugh, 26.X.1913.
- maltinii* (*Asida*), Ardoin, 1971: 47. Holotypus ♂ (133), Tunisia: La Gali-te, 2-25.VIII.1968 (A. Ceccanti, G. Adriani e M. Romiti). Allotypus ♀ (134), Tunisia: Isola Galitone, arc. Galita, 30.V.1966 (B. Lanza e S. Carfi).
- maresi* (*Akis*), Ardoin, 1971: 49. Holotypus ♂ (137), Tunisia: Isola Aguglia (= La Fauchelle) (arc. Galita), 1.VI.1966 (B. Lanza e S. Carfi). Allotypus ♀ (138), stessi dati. Paratypi (140), Tunisia: Isola Galitone, arc. Galita, 30.V.1966 (B. Lanza e S. Carfi), 2 ex.
- martellii* (*Heterocheira*), Ardoin, 1974: 143. Holotypus (2001), Somalia: Sar Uanle, 6.XI.1971 (S.B.S.-P.L.), (*). Allotypus (2002), stessi dati,

- (*). Paratypi (2003), Somalia: Sar Uanle, 20.X-14.XI.1971 (S.B.S.-P.L.), 22 ex, (*).
- martellii* (*Zophosis*), Ardoin, 1974: 138. Holotypus ♂ (1095), Somalia: Sar Uanle, 23.X.1971 (S.B.S.-P.L.), (*). Allotypus ♀ (1096), Somalia: Sar Uanle, 3.XI.1971 (S.B.S.-P.L.), (*). Paratypi (1094), Somalia: Sar Uanle, 13.X-28.XI.1971 (S.B.S.-P.L.), 33 ex, (*).
- parcepilosum* (*Gonocephalum inquinatum* ssp.), Gridelli, 1940: 240. Paratypi (7211), [Etiopia]: Eritrea, Adi-Ugri, VII.1901 (A. Andreini), 3 ex.
- rotundatum* (*Gonocephalum*), Gridelli, 1940: 241. Syntypi (7198), [Etiopia]: Eritrea, Adi-Caié, IV-X.1902 (A. Andreini), 13 ex. Syntypus (7199), [Etiopia]: Eritrea, Adi-Ugri, VII.1901 (A. Andreini). Syntypi (7200), [Etiopia]: Eritrea, Saganeiti, IV.1901 (A. Andreini), 2 ex. Syntypi (7201), [Etiopia]: Eritrea, Altip. Goaita Assaorta, V.1902 (A. Andreini), 3 ex. Syntypi (7202), [Etiopia]: Eritrea, Adigrat, 23-25.IX.1901 (A. Andreini), 2 ex.
- sardiniensis* (*Pseudoseriscus olivieri* ssp.), Leo, 1982: 102. Paratypus (7204), [Italia]: Sardegna, Cagliari, Giorgino, 15.VIII.1976 (C. Meloni).
- somalicus* (*Diphyrrhinchus*), Ardoin, 1974: 140. Holotypus ♂ (1098), Somalia: Sar Uanle, 9.XI.1971 (S.B.S.-P.L.), (*). Allotypus ♀ (1099), stes-
si dati, (*). Paratypi (2000), Somalia: Sar Uanle, 20.X-15.XI.1971 (S.B.S.-P.L.), 46 ex, (*).
- stefaninii* (*Strongylium*), Pic, 1928a: 25. Holotypus (7208), Somalia: Lugh, 26.X.1913.
- sulcitana* (*Pimelia angusticollis* ssp.), Leo & Pisano, 1978: 80. Paratypi (7205), [Italia]: Sardegna, prov. Cagliari, Porto Vesme, 20.IV.1975 (P. Leo), 2 ex.

ALLECULIDAE

- rufithorax* (*Pseudocistela*), Pic, 1928a: 24. Holotypus (7165), Somalia [ex]
italiana: Giumbo, VI.1913.

MORDELLIDAE

- andreinii* (*Mordellistena*), Pic, 1933: 126. Syntypi (7134), [Etiopia]: Eritrea, M.te Dongollo, 25.I.1901 (A. Andreini), 2 ex. Syntypus (7135), [Etiopia]: Eritrea, Adi-Ugri, VIII.1901 (A. Andreini).

- erythraeana* (*Mordellistena*), Pic, 1933: 126. Syntypus (7136), [Etiopia]: Eritrea, M.te Dongollo, 25.I.1901 (A. Andreini).
reducta (*Anaspis 4-maculata* var.), Pic, 1933: 127. Syntypi (7137), [Etiopia]: Eritrea, Adi-Caié, VIII.1902 (A. Andreini), 6 ex.

ADERIDAE

- erythreus* (*Hylophilus*), Pic, 1933: 124. Syntypus (7143), [Etiopia]: Eritrea, Adi-Caié, VIII.1902 (A. Andreini).

MELOIDAE

- andreinii* (*Zonabris*), Pic, 1933: 123. Syntypi (7133), [Etiopia]: Eritrea, Maio, Assaorta Alta, VIII.1902 (A. Andreini), 5 ex.
lanzai (*Actenodia unimaculata* ssp.), Kaszab, 1973: 51. Holotypus (670), Somalia: Giohar, 28.IV.1968 (B. Lanza e A. Simonetta), (*). Paratypus ♀ (671), stessa località, 30.IV.1968 (B. Lanza), (*).
paolii (*Epicauta testaceipes* var.), Pic, 1928b: 41. Holotypus (7131), Somalia: Villaggio Duca Abruzzi, VIII.1926 (Miss. ent. Paoli).
paolii (*Zonitoschema eborina* var.), Pic, 1928a: 23. Holotypus (7132), Somalia: Matagassile, 1913.
postmarginata (*Coryna*), Pic, 1928b: 40. Holotypus (7130), Somalia: Villaggio Duca Abruzzi, I-III.1926 (Miss. ent. Paoli).
puccionii (*Mylabris tripartita* ab.), Kaszab, 1973: 46. Holotypus (657), Somalia, 1924 (Stefanini e Puccioni).
simillima (*Cyaneolytta depressicornis* ssp.), Kaszab, 1973: 54. Holotypus (632), Somalia: Afgoi, XI.1959 (A. Sammiceli).
somalica (*Coryna*), Kaszab, 1973: 48. Holotypus ♀ (668), Somalia: Kurtum Uaro, 14.VIII.1959 (Missione Biologica in Somalia).

ANTHICIDAE

- andreinii* (*Anthicus*), Pic, 1933: 125. Holotypus (7152), [Etiopia]: Eritrea, Adi-Caié, V.1902 (A. Andreini).
andreinii (*Formicomus*), Pic, 1933: 124. Holotypus ♀ (7150), [Etiopia]: Eritrea, Adi-Ugri, VIII.1901 (A. Andreini).

- chiaromonte* (*Anthicus*), Pic, 1933: 126. Holotypus (7153), [Etiopia]: Eritrea, Adi-Caié, V.1902 (A. Andreini).
- hispidipennis* (*Anthicus*), Pic, 1933: 125. Holotypus (7151), [Etiopia]: Eritrea, Adi-Caié, VII.1902 (A. Andreini).
- paolii* (*Formicomus*), Krekich-Strassoldo, 1928: 384. Syntypi ♂♂ e ♀♀ (7154), Somalia mer.: Vill. Duca Abruzzi, 15.IX.1926 (Miss. ent. Paoli), 3 ex.

CHRYSOMELOIDEA

CERAMBICIDAE

- africanus* (*Linopteridius*), Villiers, 1972: 276. Paratypus (568), Somalia: Ola Uager, 15-18.VIII.1970 (S.B.S.), (*).
- albogrisea* (*Prosopocera octomaculata* ssp.), Breuning, 1940: 214. Holotypus (574), Somalia: Gelib, Alessandra, IV.1937 (F. Bigi).
- albovittatus* (*Parhippopsicon*), Breuning, 1978: 154. Holotypus (3617), Somalia: Dusa Mareb, 25.X.1973 (B. Lanza), (*).
- antennatus* (*Cantharocnemis*), Franz, 1938: 201. Holotypus (6910), Somalia: Obbia, Durgale-Magghiole, 1924 (Stefanini e Puccioni). Paratypi (6911), stessi dati, 2 ex.
- baldasseronii* (*Idactus*), Breuning, 1958: 107. Holotypus (578), Etiopia: Borana, Javello, XI-XII.1938-I-III.1939 (E. Faà di Bruno).
- basiruficornis* (*Exocentrus*), Breuning, 1958: 108. Holotypus (570), Somalia meridionale: Vill. Duca Abruzzi, IV-V.1926 (Miss. ent. Paoli).
- bituberosoides* (*Sophronica*), Breuning, 1972: 290. Holotypus (223), Somalia: Giohar, 18-28.IV.1968 (B. Lanza e A. Simonetta), (*). Paratypus (224), stessi dati, (*).
- brunoi* (*Lasiopezus*), Breuning, 1972: 286. Holotypus (221), Etiopia: Borana, Javello, XI-XII.1938-I-III.1939 (E. Faà di Bruno).
- distinguenda* (*Eunidia*), Aurivillius, 1928: 480. Holotypus (539), Somalia meridionale: Vill. Duca Abruzzi, IV-V.1926 (Miss. ent. Paoli) (coll. Paoli, Museo Civico di Genova).
- fuscolateralis* (*Phytoecia*), Breuning, 1977: 297. Holotypus (3627), Somalia: Sar Uanle, 24.X.1971 (S.B.S.-P.L.), (*). Paratypus (3625), stessa località, 31.X.1971, (*).
- macrophthalma* (*Eunidia*), Breuning, 1940: 217. Holotypus (575), Somalia: Gelib, Alessandra, IV.1937 (F. Bigi).

- martellii* (*Thapsyrus*), Villiers, 1972: 278. Holotypus (215), Somalia: Giohar, 18-28.IV.1968 (S.B.S.), (*).
- meleagris* (*Eunidia*), Aurivillius, 1928: 481. Holotypus (540), Somalia: Lugh, X.1913 (coll. Paoli, Museo Civico di Genova).
- metallescens* (*Sophronica*), Breuning, 1978: 153. Holotypus (3678), Somalia: Baidoa, VII.1972 (S.B.S.), (*).
- monoxenoides* (*Jordanoleiopus*), Breuning, 1972: 291. Holotypus (225), Somalia: Eggoi, Vill. Duca Abruzzi, 22-26.VIII.1962 (Miss. Biol. Somalia 62).
- postbrunnea* (*Oberea*), Breuning, 1977: 296. Holotypus (3626), Somalia: Sar Uanle, 6.XI.1971 (S.B.S.-P.L.), (*).
- quadrifasciata* (*Eunidia*), Breuning, 1940: 218. Holotypus (576), Somalia: Gelib, Alessandra, IV.1937 (F. Bigi).
- rufolimbata* (*Brunoeme*), Villiers, 1972: 275. Holotypus (214), Etiopia: Borana, Javello, XI-XII.1938-I-III.1939 (E. Faà di Bruno).
- sammichellii* (*Hololeprus*), Villiers, 1972: 280. Holotypus (217), Somalia: Afgoi, IX.1959 (A. Sammicheli).
- somalica* (*Eunidia*), Breuning, 1977: 294. Holotypus (3624), Somalia: Sar Uanle, 29.V.1973 (S.B.S.-P.L.), (*).
- somaliensis* (*Crossotus*), Breuning, 1972: 289. Holotypus (222), Somalia: Jesomma, 15.VIII.1968.
- somaliensis* (*Derolus*), Villiers, 1972: 280. Holotypus (216), Somalia: Giohar, 18-28.IV.1968 (B. Lanza e A. Simonetta), (*).
- somaliensis* (*Exocentrus*), Breuning, 1958: 108. Holotypus (577), Etiopia: Borana, Javello, XI-XII.1938-I-III.1939 (E. Faà di Bruno).
- somaliensis* (*Hyllisia*), Breuning, 1972: 284. Holotypus (219), Somalia: Giohar, 18-28.IV.1968 (B. Lanza e A. Simonetta), (*).
- somaliensis* (*Idactus*), Breuning, 1972: 286. Holotypus (220), Somalia: Dinsor, 18-19.VII.1962 (Miss. Biol. Somalia 62).
- somaliensis* (*Mimononymoides*), Breuning, 1972: 283. Holotypus (218), Somalia: Bur Dinsor, 23.VII.1962 (Miss. Biol. Somalia 62).
- somaliensis* (*Tetropiopsis*), Villiers, 1972: 273. Holotypus (213), Somalia: Bender Cassim (Migiurtinia), 24.IV.1938 (Mordini).
- x-signata* (*Paramecyna*), Breuning & Chelazzi, 1978: 175. Holotypus (3624), Zaïre: Fiume Mongo, 19-24.III.1975, (S.B.Z.), (*).

BRUCHIDAE

- arenarum* (*Caryedon*), Decelle, 1979a: 84. Holotypus ♂ (6711), Somalia: dune tra Obbia e Durghale, 1924 (Stefanini e Puccioni).

- buettikeri* (*Bruchidius*), Decelle, 1979c: 321. Paratypus (6707), Somalia meridionale: Bar-Medeghé, 1924 (Stefanini e Puccioni).
- grandis* (*Caryedon*), Decelle, 1979a: 85. Holotypus ♂ (6712), Somalia: Run 7-19.VIII.1969 (S.B.S.), (*). Allotypus ♀ (6713), Somalia: Run (Garoe), 2.VIII.1964 (Miss. Biol. Somalia 1964). Paratypi (6714, 6715), Somalia: El Bur, 5-9.IX.1962 (Miss. Biol. Somalia 1962), 2 ex.
- somalicus* (*Callosobruchus*), Decelle, 1979a: 79. Holotypus ♂ (6705), Somalia: Sar Uanle, 13.VI.1973 (S.B.S.-P.L.), (*).
- somalicus* (*Spermophagus*), Decelle, 1979a: 82. Holotypus ♂ (6710), Somalia: Sar Uanle, 30.V.1973 (S.B.S.-P.L.), (*).

CHRYSEMELIDAE

- andreinii* (*Cassida*), Spaeth, 1933: 48. Holotypus (7162), [Etiopia]: Eritrea, Cheren, IV.1903 (A. Andreini).
- andreinii* (*Cryptocephalus decoratus* var.), Pic, 1933: 129. Syntypus (7160), [Etiopia]: Eritrea, Adi-Caié, Manara, 1901 (A. Andreini).
- apicalis* (*Anteriscus*), Pic, 1933: 130. Holotypus (7158), [Etiopia]: Eritrea, Adi-Ugri, VII.1901 (A. Andreini).
- atrimembris* (*Leucastea*), Pic, 1933: 128. Holotypus (7156), [Etiopia]: Eritrea, Adi-Ugri, VIII.1901 (A. Andreini).
- chiaromontei* (*Cryptocephalus decoratus* var.), Pic, 1933: 129. Syntypus (7159), [Etiopia]: Eritrea, Adi-Caié, Manara, V.1901 (A. Andreini).
- perpusilla* (*Dactylispa*), Gestro, 1911: 28. Syntypus (7163), [Etiopia]: Eritrea, Saganeiti, IV.1901 (A. Andreini). Syntypus (7164), [Etiopia]: Eritrea, Adi-Ugri, VIII.1901 (A. Andreini).
- testaceipes* (*Crioceris*), Pic, 1933: 128. Holotypus (7157), [Etiopia]: Eritrea, Adi-Caié, IX.1902 (A. Andreini).
- ugriensis* (*Cryptocephalus decoratus* var.), Pic, 1933: 129. Syntypus (7161), [Etiopia]: Eritrea, Adi-Ugri, VII.1901 (A. Andreini).

CURCULIONOIDEA

ANTHRIBIDAE

- targionii* (*Cratoparis*), Piccioli, 1877: 214. Holotypus (7364), [Italia: Toscana], dintorni di Firenze (Porte Sante).
- ricini* (*Misthosimella*), Jordan, 1929: 165. Holotypus ♀ (7363), Somalia mer.: Villaggio Duca Abruzzi (Miss. ent. Paoli).

BRENTHIDAE

- adustus* (*Cerobates*), Senna, 1894b: 184. Syntypi (6937), [Indonesia]: Java Occ., Toegoe (J. D. Pasteur), 10 ex.
- aeneus* (*Ubanius*), Senna, 1895a: 222. Syntypus ♀ (7004), Panama: Chiriqui, Amer. Centr. (don. Standinger).
- amazonicus* (*Raphidorrhynchus*), Senna, 1894e: 598. Holotypus ♀ (7019), Brasile: Amazonas (don. Standinger).
- ambiguus* (*Diurus*), Senna, 1909: 37. Holotypus ♂ (7050), [Sarawak]: N.W. Borneo, Kuching, 25.I.1900 (by Dyak coll., pres. 1900 by Shelford). Allotypus ♀ (7051), stessa località, 1.XI.1899.
- amplicollis* (*Raphidorrhynchus*), 1894e: 597. Holotypus ♀ (7020), Brasile: Fonteboa (d. Standinger).
- andamanicus* (*Cerobates*), Senna, 1896c: 223. Holotypus (6943), [India]: Is. Andaman (da E. André).
- andrewsi* (*Caenorychodes*), Gahan, 1900: 117. Paratypi (7009), [Australia]: Christmas I., IV.1898 (C.W. Andrews), 4 ex.
- angulaticeps* (*Jonthocerus*), 1898a: 53. Syntypi ♂♂ e ♀♀ (6925), [Indonesia]: Java occ. (J.D. Pasteur), 7 ex.
- angusticeps* (*Cyphagogus*), Senna, 1893c: 296. Holotypus ♂ (6913), [Indonesia]: Sumatra, sul tabacco secco (A. Grouvelle).
- angustipennis* (*Cerobates*), Senna, 1894b: 182. Syntypi (6936), [Indonesia]: Java occ., Toegoe (J.D. Pasteur), 9 ex.
- armaticeps* (*Cordus*), Senna, 1894a: 559. Syntypus (6977), Nuova Guinea.
- articulatus* (*Diurus*), Senna, 1909: 38. Holotypus ♂ (7047), Sarawak: N.W. Borneo, Kuching, 31.X.1899 (by Dyak coll., pres. 1900 by D.R. Shelford).
- aureopilosus* (*Trachelizus*), Senna, 1889b: 105. Holotypus ♀ (6960), [Brasile]: Fonteboa (don. Standinger).
- australiana* (*Miolispa*), Senna, 1897a: 58. Holotypus ♂ (6963), Australia: Victoria (ex Desbrochers). Allotypus ♀ (6964), stessi dati.
- australicus* (*Araiorrhinus*), Senna, 1893c: 327. Holotypus (6948), Australia: New S. Wales (don. Faust).
- bicolor* (*Bolbocranius*), Senna, 1898c: 371. Syntypi (7070), N. Camerun: Station J. Albrechts Höhe (Conradt), 8 ex.
- birmanicus* (*Cerobates*), Senna, 1894c: 364. Syntypi (6940), N. Birmania: Rubis Mines (H. Doherty), 2 ex.
- birmanicus* (*Miolispoides*), Senna, 1894c: 370. Holotypus ♂ (7000), Haute Birmanie: Mines d. Rubis (W. Doherty).

- borneensis* (*Miolispa*), Senna, 1898b: 70. Holotypus ♂ (6965), Borneo (ex coll. Desbrochers).
- brevisulcatus* (*Microtrachelizus*), Senna, 1894a: 558. Holotypus (6950), N. Guinea: Ighibirei, VII-VIII.1890 (L. Loria).
- cardoni* (*Symmorphocerus*), Senna, 1894d: 385. Syntypi ♀ ♀ (6976), [India]: Barway (P. Cardon), 4 ex.
- carinensis* (*Jonthocerus*), Senna, 1894c: 362. Syntypus ♀ (6926), [Birmaniam]: Carin Asciiui-Ghecù, 1400-1500 m, III-IV.1888 (L. Fea). Syntypus ♀ (6927), [Birmaniam]: Carin Chebà, 900-1000 m, V-XII.1888 (L. Fea). Syntypi (6995), Alta Birmania: Miniere di Rubini, 1200-2300 m, 2 ex.
- carinirostris* (*Schizotrachelus*), Senna, 1892e: 489. Syntypus ♂ (7066), Birmania: Bharno, VII.1886 (L. Fea).
- celebicus* (*Heteroclites*), Senna, 1897b: 242. Syntypi (7042), [Indonesia]: S. Celebes, Lompa-Battau, 3000, III.1896 (H. Fruhstorfer), 5 ex. Syntypi (7043), [Indonesia]: S. Celebes, Bua-Kraeng, 5000, II.1896 (H. Fruhstorfer), 2 ex.
- ceylonicus* (*Euphenges*), Calabresi, 1920b: 63. Holotypus ♂ (7015), Ceylon (A. Boucard).
- chiriquensis* (*Tmetogonus*), Senna, 1895a: 225. Holotypus (7008), Amer. centr. [Panama]: Chiriqui (don. Standinger).
- cinereus* (*Ulocerus*), Senna, 1896a: 208. Syntypus ♀ (7059), Brasile (d. Mus. Tring).
- cocoensis* (*Raphidorrhynchus*), Calabresi, 1920b: 71. Holotypus ♂ (7021), Costa Rica: Is. de Coco, VI.1898.
- compressitarsis* (*Apterorhinus*), Senna, 1896b: 59. Holotypus ♂ (7071), [Filippine]: S. Luzon, Albay, 1894 (Whitehead).
- compressithorax* (*Microsebus*), Senna, 1898c: 370. Holotypus (6918), S.W. Cameroun: Stat. Lolodorf (Conradt).
- conradti* (*Jonthocerus*), Senna, 1898c: 374. Holotypus ♂ (6932), N. Cameroun: Stat. J. Albrechts Höhe (Conradt).
- consanguineus* (*Raphirhynchus*), Senna, 1895a: 217. Syntypi (7022), Venezuela: Merida (Tring Mus.), 4 ex.
- contiguus* (*Trachelizus*), Senna, 1893a: 266. Holotypus (6952), [Indonesia]: Engano, Malaconni, VI.1891 (E. Modigliani).
- costipennis* (*Acramorphocephalus*), Calabresi, 1920c: 66. Syntypus ♂ (6981), N. Cameroun: Stat. J. Albrechts Höhe (Conradt). Syntypus ♀ (6982), Congo Franç. [Rep. Dem. Congo]: Bènitò.
- costipennis* (*Oncodemerus*), Senna, 1892c: 157. Holotypus ♂ (6917), Gabon (da Standinger).

- crassus* (*Pseudorychodes*), Senna, 1894c: 378. Paratypus (7013), Alta Birmania: Min.e rubini (H. Doherty).
- cruciata* (*Miolispa*), Senna, 1898b: 69. Holotypus ♂ (6966), N. Borneo (ex coll. Fruhstorfer).
- cruentatus* (*Cerobates*), Senna, 1898c: 372. Holotypus ♂ (7088), N. Camerun: Stat. Joh. Albrechts Höhe (Conradt).
- discors* (*Miolispa*), Senna, 1895b: 358. Holotypus ♀ (6967), [Malesia]: Penang [= Pinang] (ex A. Boucard).
- emmae* (*Hopliterrhynchus*), Senna, 1892b: 26. Holotypus ♂ (7032), Malesia (A. Boucard).
- enganica* (*Miolispa*), Senna, 1893a: 267. Paratypus ♂ (7001), [Indonesia]: Engano, Bua-Bua, V-VI.1891 (E. Modigliani).
- exportatus* (*Araiorrhinus*), Senna, 1893c: 327. Holotypus (6946), [Indonesia]: Sumatra, imp. c. tabacco (da Grouvelle).
- fausti* (*Araiorrhinus*), Senna, 1893c: 325. Syntypi ♂ ♂ (6957), S.O. Kamerun: Stat. Lolodorf (H. Conradt), 2 ex.
- feae* (*Callipareius*), Senna, 1892e: 445. Paratypus ♂ (6919), Birmania: Carin [montagne], Ghecù [distretto], 1300—1400 m, V.1888 (L. Fea).
- festivus* (*Brenthus*), Senna, 1897a: 232. Holotypus ♂ (7080), Columbia: Canca (d. Standinger). Paratypi (7082), Venezuela: Merida, 12 ex. Paratypi (7083), Panama: Chiriqui, 2 ex. Paratypi (7081), Columbia (A. Boucard), 5 ex.
- filicauda* (*Diurus*), Senna, 1909: 47. Holotypus ♂ (7055), Sarawak: Borneo (da Standinger). Allotypus ♀ (7056), [Sarawak]: N.W. Borneo: Kuching capt., 25.I.1900 (by Dyak coll., pres. 1900 by Shelford).
- flavornatus* (*Cephalobarus macrocephalus* var.), Senna, 1889a: 35. Syntypi ♂ e ♀ (7073), Columbia: [rio] Canca (d. Standinger), 2 ex.
- forcipitigerus* (*Arrhenodes*), Gyllhenhal, 1840: 478. Paratypus ♂ (7078), Cuba (A. Boucard).
- foveatus* (*Callipareius*), Senna, 1894d: 384. Paratypus (6920), [India]: Kur-seong (P. Kraet).
- foveolatus* (*Jonthocerus*), Senna, 1893c: 300. Lectotypus ♂ (6930), [Indonesia]: Sumatra, imp. in Europa col tabacco secco (Grouvelle) [designato da Damoiseau, *in litteris*].
- fruhstorferi* (*Prodeutor*), Senna, 1897b: 241. Syntypi (7044), [Indonesia]: S. Celebes, Bua-Kraing, 5000, II.1896 (H. Fruhstorfer), 8 ex.
- fruhstorferi* (*Pseudorychodes*), Senna, 1894c: 375. Allotypus ♀ (7011), [Indonesia]: Java occ., m. Tjikorae, 4000, 1892 (H. Fruhstorfer).
- funebis* (*Arrhenodes*), Sharp, 1895: 44. Syntypi ♂ ♂ e ♀ ♀ (7002), Panama: V. de Chiriqui, 25-4000 ft (Champion), 4 ex.

- fusca* (*Heteroblysmia*), Calabresi, 1920c: 66. Holotypus ♂ (7017), [Sarawak]: N.W. Borneo, Kuching, 11.VII.1900 (by Dyak coll., pres. 1900 by Shelford).
- gestri* (*Pithoderes*), Calabresi, 1920d: 43. Holotypus ♂ (7067), [Rep. Pop. Congo]: Bènit.
- gestroi* (*Stereodermus*), 1893a: 261. Syntypi (6934), [Indonesia]: Engano, Bua-Bua, V-VI.1891 (E. Modigliani), 3 ex.
- gestroi* (*Uropterus*), Senna, 1894a: 562. Syntypi (7060), N. Guinea: Ighibirei, VII-VIII.1890 (L. Loria), 3 ex.
- grouvellei* (*Cerobates*), Senna, 1893c: 307. Lectotypus ♀ (6938), [Indonesia]: Sumatra, imp. c. tabacco (da Grouvelle) [designato da Damoiseau, in litteris].
- grouvellei* (*Higonius*), Senna, 1893c: 311. Holotypus (6944), [Indonesia]: Sumatra, imp. c. tabacco (da Grouvelle).
- guatemalensis* (*Nemocephalus*), Senna, 1893b: 128. Syntypi (7069), Guatemala (A. Boucard), 2 ex.
- guatemalensis* (*Claedoderes*), Sharp, 1895: 52. Syntypus ♂ (7074), Guatemala: Capetillo (G.C. Champion). Syntypus ♀ (7124), Guatemala: Cerro Zunil, 4-5000 ft (Champion).
- hirsutus* (*Hyperephanus*), Senna, 1892c: 155. Holotypus (6933), Cile (ex coll. Boucard).
- hybridus* (*Cerobates*), Senna, 1898c: 371. Holotypus (6939), N. Kamerun: Station Albrechts Höhe (L. Conradt).
- insculptus* (*Raphidorrhynchus*), Senna, 1889a: 36. Holotypus ♀ (7023), Brasile: Fonteboa (d. Standinger).
- insularis* (*Trachelizus*), Senna, 1892a: 163. Holotypus ♀ (6961), [Malesia]: Is. Penang [= Pinang], (A. Boucard). Allotypus ♂ (6962), [Indonesia]: Giava (A. Boucard).
- intermedia* (*Miolispa*), Senna, 1897b: 239. Holotypus ♂ (6984), [Indonesia]: Nord Celebes, Toli-Toli, XI-XII.1895 (H. Fruhstorfer).
- intermedius* (*Diurus*), Senna, 1909: 41. Holotypus (7049), [Sarawak]: N.W. Borneo, Kuching. capt., 20.III.1900 (by Dyak coll., pres. 1900 by R. Shelford).
- intermedius* (*Schizotrachelus*), Senna, 1892e: 481. Syntypi (7061), Indonesia: Java occ. (W. Pasteur).
- interruptus* (*Schizotrachelus*), Senna, 1898b: 72. Holotypus ♂ (7065), [Malesia]: Is. Penang [= Pinang] (d. Standinger).
- iridescent* (*Euphenges*), Calabresi, 1920b: 65. Syntypi ♂ e ♀ (7016), [Malesia]: Is. Penang [= Pinang] (A. Boucard), 2 ex.

- javanica* (*Miolispa*), Senna, 1892a: 167. Paratypi (6990), [Indonesia]: Java occ. (W. Pasteur), 27 ex.
- jordani* (*Miolispa*), Senna, 1895b: 360. Holotypus ♂ (6983), [Malesia]: Is. Penang [= Pinang] (ex. Boucard).
- kleinei* (*Anepsiotus*), Calabresi, 1920b: 68. Paratypi (7036), [Malesia]: Is. Penang [= Pinang] (A. Boucard), 2 ex.
- kraatzi* (*Cordus*), Senna, 1898c: 375. Holotypus ♂ (6978), S.O. Camerun: Station Lolodorf (L. Conradt).
- laevigatus* (*Trachelizus*), Senna, 1898a: 58. Holotypus (7086), [Indonesia]: Java occ., 1895 (W. Pasteur).
- laevis* (*Usambius conradti* var.), Senna, 1898: 370. Syntypi (6915), N. Kamerun: Station J. Albrechts Höhe (L. Conradt), 2 ex. Syntypi (6916), stessa località (ex Dr. Kraatz), 2 ex.
- laevipennis* (*Cerobates*), Senna, 1896c: 219. Syntypus ♂ (6935), [Indonesia]: Is. Sanghi [= Sangis].
- lamellipes* (*Fontebuanus*), Senna, 1893b: 125. Holotypus ♂ (7068), Brasile: Fonteboa, Amazon. Fl., 1889 (Standinger).
- linearis* (*Raphidorrhynchus*), Senna, 1894e: 607. Holotypus ♀ (7024), Messico: Sierra de Durango (d. A. Boucard).
- lineata* (*Miolispa*), Senna, 1898a: 57. Paratypi ♂ ♂ (6985), [Indonesia]: Java occ., 1895 (W. Pasteur), 4 ex.
- luzonicus* (*Anepsiotus*), Calabresi, 1920b: 66. Holotypus ♂ (7038), Filippine: N. Luzon, 1894 (Whitehead).
- maculipennis* (*Brenthus*), Senna, 1897a: 234. Holotypus ♂ (7079), Ecuador (A. Boucard).
- marginatus* (*Schizotrachelus*), Senna, 1898b: 75. Holotypus ♂ (7063), Nord Borneo (ex coll. Fruhstorfer). Allotypus ♀ (7064), stessi dati.
- mariae* (*Miolispa*), Senna, 1891: 165. Holotypus ♂ (6988), [Malesia]: Is. Penang [= Pinang] (da A. Boucard). Paratypus ♂ (6989), stessi dati.
- major* (*Stratiorrhina*), Calabresi, 1921a: 47. Syntypi (7005), Borneo (A. Boucard), 2 ex. Syntypus (7006), N.W. Borneo, Kuching capt., 17.VII.1900 (by Dyak coll. on flowering tree, pres. 1900 by Shelford). Syntypus (7007), Sumatra (Dohrn) (don. Dr. Faust).
- metallica* (*Miolispa*), Senna, 1892a: 169. Syntypi (6994), [Indonesia]: Java occ., 1895 (W. Pasteur), 26 ex.
- metallicus* (*Schizotrachelus*), Senna, 1897b: 240. Syntypi (7062), [Indonesia]: S. Celebes, Lompa-Battau, 3000, III.1896 (H. Fruhstorfer), 2 ex.
- modestus* (*Trachelizus*), Senna, 1892a: 165. Syntypus (7087), [Indonesia]: Java (da F.P. Pascoe).

- modiglianii* (*Cyphagopus*), Senna, 1893a: 258. Paratypi ♂ e ♀ (6912), [Indonesia]: Engano, Bua-Bua, 5.VI.1891 (E. Modigliani), 2 ex.
- mutabilis* (*Prophthalmus*), Senna, 1893a: 274. Syntypi (6971), [Indonesia]: Engano, Bua-Bua, V-VI.1891 (E. Modigliani), 13 ex.
- niassicus* (*Higonius*), Senna, 1893c: 310. Holotypus ♂ (6945), [Indonesia]: Is. Nias (Jordan) (ex coll. Faust).
- nitidicollis* (*Anepsiotes*), Calabresi, 1920b: 69. Holotypus ♂ (7037), Philippines: Manila (Whitehead).
- nitidissimus* (*Proephebocerus*), Calabresi, 1920a: 39. Holotypus ♂ (6958), Madagascar: Andrangoloaka, 1600 m O.S.O. de Tananarive. Allotypus ♀ (6959), Madagascar: S de la Baie de Antongil.
- nonfriedi* (*Miolispa*), Senna, 1895b: 363. Holotypus ♂ (6986), Nuova Guinea (coll. Nonfried). Allotypus ♀ (6987), stessi dati.
- nupta* (*Miolispa*), Senna, 1892a: 171. Syntypi (7085), [Indonesia]: Java occ. (J.D. Pasteur), 20 ex.
- oberthueri* (*Hoplopisthius*), Senna, 1892d: 255. Syntypus ♂ (6954), Birmania: Min.a di rubini, Alta Birmania, m 1200-2300 (Doherty).
- orientalis* (*Pseudobelopherus*), Calabresi, 1920b: 61. Holotypus ♂ (7034), Malacca (Roelofs). Allotypus ♀ (7035), Malacca (A. Boucard).
- ornatus* (*Heterorrhynchus*), Calabresi, 1921b: 63. Holotypus ♂ (7033), Borneo, 1894 (M. Vallier).
- panamensis* (*Raphidorrhynchus*), Senna, 1894e: 609. Holotypus ♀ (7025), Panama (A. Boucard).
- pasteuri* (*Jonthocerus*), Senna, 1898a: 55. Holotypus ♂ (6928), [Indonesia]: Java occ. (J.D. Pasteur). Paratypus (6929), stessi dati.
- pasteuri* (*Miolispa*), Senna, 1898a: 56. Paratypus ♂ (6974), [Indonesia]: Java occ. (D. Pasteur).
- philippinus* (*Diurus*), Senna, 1909: 45. Syntypi (7054), [Filippine]: N. Luzon, 1894 (Whitehead), 10 ex.
- poultoni* (*Diurus*), Senna, 1909: 44. Syntypi (7053), Sarawak: N.W. Borneo, Kuching capt., 13.VII.1899 (by Dyak coll., pres. 1900 by Shelford), 2 ex. Syntypi (7054), stessa località, 12.XII.1899, 2 ex.
- propinquus* (*Acratus*), Senna, 1889b: 108. Holotypus ♂ (7075), Brasile: Fonteboa (d. Standinger). Allotypus ♀ (7076), stessi dati.
- pubescens* (*Microtrachelizus*), Senna, 1893c: 320. Holotypus (6949), [Indonesia]: Sumatra, s. tabacco (A. Grouvelle).
- punctulatus* (*Cerobates*), Senna, 1898c: 373. Holotypus ♂ (6941), N. Kamerun: Station J. Albrechts Höhe (L. Conradt).
- pygmoea* (*Miolispa*), Senna, 1895b: 360. Allotypus ♀ (6991), [Malesia]: